

QSO hosts and environments at $z = 0.9$ to 4.2 : *JHK* images with Adaptive Optics

J.B. Hutchings¹, David Crampton¹, S.L. Morris¹, and D. Durand
 Dominion Astrophysical Observatory, National Research Council of Canada,
 Victoria, B.C. V8X 4M6, Canada

E. Steinbring¹
 Dept of Physics and Astronomy, University of Victoria

ABSTRACT

We have observed nine QSOs with redshifts 0.85 to 4.16 at near-IR wavelengths with *Pueo*, the adaptive optics bonnette of the Canada-France-Hawaii telescope. Exposure times ranged from 1500 to 24000s (mostly near 7000s) in *J*, *H*, or *K* bands, with pixels $0''.035$ on the sky. The FWHM of the co-added images at the location of the quasars are typically $0''.16$. Including another QSO published previously, we find associated QSO structure in at least eight of ten objects, including the QSO at $z = 4.16$. The structures seen in all cases include long faint features which appear to be tidal tails. In four cases we have also resolved the QSO host galaxy, but find them to be smooth and symmetrical: future PSF removal may expand this result. Including one object previously reported, of the nine objects with more extended structure, five are radio-loud, and all but one of these appear to be in a dense small group of compact galaxy companions. The radio-quiet objects do not occupy the same dense environments, as seen in the NIR. In this small sample we do not find any apparent trends of these properties with redshift, over the range $0.8 < z < 2.4$. The colors of the host galaxies and companions are consistent with young stellar populations at the QSO redshift. Our observations suggest that adaptive optic observations in the visible region will exhibit luminous signatures of the substantial star-formation activity that must be occurring.

Subject headings: quasars: general, galaxies: interactions, instrumentation: adaptive optics

¹Visiting Astronomers, Canada-France-Hawaii Telescope, operated by the National Research Council of Canada, the Centre de la Recherche Scientifique de France, and the University of Hawaii

1. Introduction

QSO host galaxies are well investigated at redshifts up to ~ 0.6 , with both ground-based and *HST* images, both at visible and near infrared wavelengths (e.g. Hutchings & Neff 1992, 1997; McLeod & Rieke 1995; Bahcall et al. 1997 and references therein). Much has been written on the general evidence for tidal events being the trigger for nuclear activity, the hybrid morphology of the QSO host galaxies, associated star-formation, and the different environments of companion galaxies between radio-loud and radio-quiet QSOs.

At higher redshift, there have been ground-based data that resolve QSO hosts at redshifts in the range 2.0 to 2.5. These, particularly the radio-loud ones, appear to be very luminous galaxies, with very high star-formation rates, and with no counterparts in the present day universe (e.g. Heckman et al. 1991; Lehnert et al. 1992; Hutchings and Neff 1997b; Hutchings 1998; Aretxaga et al. 1995,8). There are also indications that QSOs of all kinds occur in small dense groups of companion galaxies also with high star-formation rates (Hutchings 1995,8; Campos et al 1998). There is a clear similarity to radio galaxies at comparable redshifts; they also are very luminous and are located in dense environments (e.g., Rottgering 1997, Pentericci et al 1998)

The high redshift QSOs observed so far lack the spatial detail and the statistical samples that are available for the lower redshift QSOs, so several key questions cannot yet be answered. Primarily these concern the cosmic evolution of the QSO phenomenon. We do not know how the nature of QSO activity evolves with redshift: do the sites of QSO activity evolve as different environments become favourable; does the environment evolve at all; or does the environment not matter? We also do not know if the trigger of nuclear activity changes with redshift (as the host galaxies age or as the interaction rate changes), or if the luminosity and visibility of QSO activity evolves. Is star-formation a cause, effect, or parallel process in QSO events? Do radio-loud and radio-quiet QSOs evolve differently?

We have begun a program to obtain high resolution and high signal-to-noise images of QSOs in the redshift range $1 < z < 4$. The data reported here are from near infrared images obtained with *Pueo*, the Adaptive Optics Bonnette of the CFHT (Rigaut et al. 1998). We have previously reported similar observations for the $z = 1.1$ optically-selected QSO 1055+019 (Hutchings et al. 1998). In this paper we add 9 more to the sample, and discuss the individual cases and the trends we see in the sample so far. As noted in Hutchings et al 1998, our observing list is restricted to QSOs with nearby bright guide stars for the AO system. Thus, while we are investigating objects with redshifts larger than ~ 0.8 , the sample available at any date of observation is not large, and our principal criteria have been to attempt to match radio-loud and radio-quiet objects, and to cover a wide range of redshift.

2. Observations and data

The data reported here were obtained with *Pueo* in January and March 1998. Table 1 gives the list of objects and Table 2 the journal of observations. All observations used a nearby guide star, as listed in the table, and corrected images at the location of the QSOs generally have $\text{FWHM} \sim 0''.2$ or better. The camera used is KIR, a 1024×1024 pixel Hg:Cd:Te detector described in the CFHT website. This detector has lower noise and less image retention than the 256 pixel MONICA camera used previously (i.e. fewer reads are required to remove the effects of bright objects in the detector). Pixels are $0''.035$, giving a field of $36''$, so that the guide star is usually in the science image as well as the QSO and its local companions. The observations were 5 minute integrations performed in a non-repeating dither pattern covering several arcsec. Conditions were good for both the runs. The uncorrected seeing was in the range $0''.5$ to $0''.9$ FWHM, and the transparency was good for all observations reported. All observations were made in standard J, H, K filters. Performance in the K band is somewhat compromised by the additional thermal background from the AO system, but unfortunately no K' or K_S filter was available for our observations. Near-infrared photometric standards (Casali and Hawarden 1992) were observed, and images were also obtained of crowded stellar fields (M91 and M15) to calibrate the variation of the PSF with distance and brightness of the guide star.

Table 2 summarizes the data. The FWHM were measured at the QSO location (these are, of course, somewhat worse than at the guide star). In the best data the FWHM typically vary from $0''.13$ at K to $0''.15$ at H and $0''.20$ at J . There were substantial differences during and between nights and, on average, the results were better during the March run than the January run.

The data were processed using ‘standard’ near infrared techniques, but in several cases independently and differently by more than one author. We describe two approaches. First, sky frames were obtained as medians of the unshifted dither frames. In cases where the saturated guide star is in the image, these were edited out before the median was computed, as otherwise they usually leave a faint imprint. In most cases we were able to use sky frames from the entire period of observation of an object. However, during nights when the sky brightness was very variable, running medians of data taken closely in time were used. After sky-subtraction the data frames were shifted to superpose the QSOs exactly, and combined with sigma clipping. (The shifts were done with linear interpolation, after checking that more elaborate shifts produced the same result.) Although flat field images of dome lights were obtained, it was found empirically that flat field division did little to improve the sky flatness or noise, presumably as the sky frames constitute a flat field, and so they were generally not used.

The small pixels and the bright sky make it hard to see faint extended features such as tidal tails or the outer parts of galaxies. Further, no very sharp morphological features were detected in any of the QSOs or their companions. Thus, to improve detection of faint features, we further smoothed the sky by editing out the QSO and companions and fitting a spline surface to the remaining sky. This was subtracted from the image to remove sky brightness variation on scales of several arcsec or more. This improved our ability to investigate and measure faint tails and flux extensions over scales up to a few arcsec. *The above process was repeated several times with slightly different edits and surface fits, and the final result was a mean of the individual results, to remove spurious artifacts.* The features we report here are seen in all processed images even before this flattening procedure, so we have good confidence in their reality. Finally, in most cases we found that images with 4×4 block-averaged pixels ($0''.14$ pixels), were able to reveal the faint extended features more clearly.

A similar procedure was used developed and used independently by one of us, to deal with changing sky brightness and the faint haloes around the bright guide stars. First, a median sky frame was generated from all target exposures on each QSO for each filter. After correction with this first-order flat-field, the exposures were grouped and combined according to mean sky-brightness, and a mask was generated which included all bright pixels. Second-order flat-fields were then generated using the masks and applied for each group. Finally, the frames were registered and median-combined.

The features we discuss are faint and difficult to display, and in some cases are comparable with sky features. However, we note that they are seen independently of the different authors' processing, often appear in more than one filter, are robust to sky smoothing processing, and have flux measures that agree from differently processed images. Features that did not pass all these criteria were rejected.

Flux measurements were made by summing the signal in pixels that include the structures of interest, and subtracting the mean nearby sky level. This was done on both the full resolution and binned data. All galaxies and symmetrical features were also measured by using the IRAF task `imexam`. Limiting surface brightnesses vary with filter and integration time, as well as sky conditions, but our rough surface brightness limits in mag per square arcsec are 22.4 in K , 23.4 in H , and 24.4 in J .

We describe below the individual QSOs and their environments, and end with some collective remarks on the ensemble. The tables for each object field give all measures made in this way. Generally, the magnitude errors for smooth round objects are ± 0.1 mag for $J = 19.0$, $H = 18.4$, $K = 17.9$, and these increase to ± 0.2 mag for objects 2.3 mag fainter and ± 0.4 mag for objects yet another 1 mag fainter. Errors in extended features are comparable,

while those for PSF-subtracted fluxes are discussed individually. In this paper we have not attempted to do careful PSF modelling and removal or deconvolution. That will be dealt with in a separate paper. We mention the results of simple PSF subtraction in a couple of cases where the QSO is sufficiently well resolved that this is not a significant issue.

We quote absolute magnitudes at the QSO redshift, for objects that have the size, brightness, and colors to be possible companions. These are for an assumed $H_0 = 75 \text{ km s}^{-1} \text{ kpc}^{-1}$ and $q_0 = 0.5$, with no K-correction (K-corrections depend on stellar population and bandpass, but for young populations at these redshifts they are likely to be close to zero: -1 mag for B0 stars and zero for A0 stars). Comparison of these luminosities indicates that the possible companions (and QSO host galaxies) have rest wavelength values in the range typical of bright galaxies. If they have active star-formation, they will be more luminous at shorter rest wavelengths. The shortest rest wavelengths we are observing range from $4600 \text{ \AA}$ to $7000 \text{ \AA}$, for the three *J* band observations. In *H* band, these range from $4900 \text{ \AA}$ to $9000 \text{ \AA}$.

Figure 1 shows the distribution of detected objects in all the fields. Measures of most of them are given in individual tables. The figure indicates which objects are assumed from their size and/or brightness to be foreground galaxies (F), and stars (*), as well as the guide stars when they appeared in the field. We discuss them below in order of increasing redshift.

3. 0915–213

This is a radio-loud QSO at $z = 0.85$. It has a flat radio spectrum, of medium flux (0.6 Jy at 5 GHz), and is probably compact, but there is no published radio map. Our data are in the *H* and *K* bands. In both, the QSO lies in a group of four faint objects that occupy one quadrant of the KIR image - the rest of the field is empty, with the exception of two larger low-surface-brightness galaxies, (and two stars) which are widely spaced. We suppose these to be foreground galaxies with low surface brightness.

The QSO is extended asymmetrically to the NE, and lies very close ($1''.4$) to the faintest compact companion (#3), to the NW. A fainter companion (#1) lies further to the N (see Figure 2). There is an extended wisp of luminosity beyond the line to the nearby companion, which turns through 90° and brightens slightly, some $4''$ from the QSO. These are very faint features but they show up on subsets of our data and also in both colors, so we regard them as real. Table 3 shows photometric measures of objects in the field of 0915–213.

We have not yet obtained good PSF subtraction in the central QSO, either because of

real complex structure, or because the PSF distortion away from the guide star is unusually large. However, preliminary results indicate that the QSO host is elliptical with its long axis in the direction of the large extended luminosity to the NE. The fluxes given for the QSO host are derived from the subtractions and also from the azimuthally averaged profiles, which agreed to within 30%.

The host galaxy extension differs between H and K , suggesting a color gradient. The nearby companion does not have a bright nucleus and is extended tangentially to the direction to the QSO. While signal levels are low, we see no structure within the features described, and their overall structure is very suggestive of tidal extension of an old population of stars associated with the QSO host galaxy (see e.g. 2141+175 in Hutchings et al 1994) .

4. 0804+499

This QSO, at $z = 1.43$, was observed in H and J bands. It is a radio source that is unresolved at subarcsecond levels. The field is quite empty, containing only the guide star, another object comparable with the QSO, and a single faint galaxy some $20''$ away in both H and J band. In J band, we also see a very faint galaxy $10''$ to the S of the QSO. Thus, if there are companions, they are faint in the near infrared. Table 4 shows our photometry of the objects.

The QSO does show a faint straight ‘jet’ or tail in both J and H , extending $3''.5$ to the N (see figure 3). The feature is almost parallel to the diffraction spikes seen in the guide star, but is at 11° to it, and emerges off-centered from the QSO. Thus, we consider it is real, but it is difficult to measure accurately because of its faintness, the varying sky near the guide star, and the presence of diffraction spikes. Measures were made after surface fitting and subtraction of the sky, and also after rotation and subtraction, both in full resolution and 4×4 block-averaged images.

The jet is relatively featureless, but appears to have a small opening angle. It suggests, once again, that there has been a recent tidal event, even in this apparently empty field. Deep optical images may show if there are companions with younger stellar populations present. The color of the jet is comparable with that of the whole QSO, but its value is poorly determined.

Aside from the jet, the QSO image is not detectably extended, and has the same profile as the stars. However, the image resolution is lower than average in these data ($0''.24$) due to the fainter guide star.

5. 1337–013

This is a radio-loud QSO at $z = 1.62$. The radio source is compact and variable, with no published structure. Our data show the QSO to have at least 4 companions within $15''$. The closest one marked in Figure 1 lies about $4''$ to the NE. However, there is also a resolved faint knot $2''$ S of the QSO (see figure 4) and possible faint flux extending to it. These nearby features are seen more clearly in J band, and they have remarkably blue colors. Table 5 shows the measurements on the objects.

This QSO is another that appears to be in a group of galaxies, and interacting with one of its close neighbours. The blue color of the close companions suggests that they are both dominated by active star-formation, presumably as result of the tidal events with the QSO, or its nuclear radiation: they are projected at 10 and 20 kpc from the QSO nucleus.

6. 1540+180

This is a radio-loud quasar at $z = 1.66$ (4C18.43) listed with approximate visual magnitude 18 by Hewett & Burbidge (1993). The radio structure is a bent FR II structure with bright knots along the lobes, which suggests that there is interaction with dense and clumpy IGM. We obtained images at J , H , and K , with good signal over approximately $12''$ radius around the QSO. In these images (Figure 5 shows the region near the QSO) the quasar is the brightest object, lying in a uniformly populated region of eight galaxies which are up to 3 mag fainter than the QSO itself. It is possible that these are members of a group associated with the quasar, from their high density and similar colors. Table 6 shows photometry of several: the others lie too close to the edge of the field to enable good measures.

Figure 5 shows the vicinity of the QSO. The closest object (to the E) seen clearly in J and H is an artifact produced by the guide star: there are several of these in a characteristic pattern. The K band image shows a clear NE extension to the QSO and possible faint luminosity further to the N. The H band image does not show the bright arm to the NE. However, there is a curved region of extended luminosity extending about $4''$ to the N and ending in a brighter region. In the J band image this long curved region is only marginally detected, but the bright region at the end is present. This luminosity is in the general direction of the nearest galaxy but does not connect with it, to the limit of our detection. We see no luminosity to the S of the QSO.

The radio structure to the N matches the near infrared faint luminosity quite closely, but not exactly. The radio lobe to the N has the same length and lies mostly alongside,

to the E of the near infrared flux, and the edge-brightened lobe ends next to the brighter region of H and J emission. The radio structure suggests that the N lobe is the more active at the present time. The K band extension of the QSO does not match the direction of the radio structure nearest the nucleus.

This is the only field for which we have three colors, and are able to make a color-color plot to compare with stellar population models. Two-color photometry of the QSO and its companions is shown in figure 6, compared with age tracks for stellar populations at different redshifts. Object 5 appears as a larger disk galaxy and in this plot corresponds with a slightly reddened galaxy of intermediate age at redshift ~ 0.3 . The other nearby galaxies lie in positions on the plot that correspond to the QSO redshift. Another compact galaxy near the edge of the field is not seen in all colors due to pointing offsets, but may also be a companion. The QSO nucleus has the bluest color, but the colors of the companions indicate they are all young, with a spread of age or dust content. The objects are compact, with diameters in our image from 2 to 5 kpc (at 5 kpc per arcsec: all are smaller than one arcsec). Thus, this appears to be a compact group of galaxies dominated by young stellar populations. A similar situation is seen around the $z = 2.2$ QSO 1345+580 (Hutchings 1998). The galaxies are luminous and comparable with that of the QSO host, and lie within a diameter of less than 100 kpc: thus this is a very dense group, which is likely to evolve by merging or breaking up over cosmic time.

The QSO has several lower redshift absorption line systems, around $z=0.7$, and one at $z=1.46$. The companion galaxy colors do not correspond to objects at $z=0.7$, but we cannot exclude their lying at 1.46, which is close to the emission line redshift of 1.66.

Crude PSF subtraction in K band shows the NE tail clearly and indicates a resolved flux comparable to the companion galaxies (see figure 7). The QSO is well resolved in this band and the subtracted host galaxy flux is not significantly dependent on the PSF correction for distance from the guide star. In H band, however, the resolution of the host galaxy is marginal and depends strongly on the PSF correction. In the K band resolved flux, photometric K -corrections are likely to be no more than a few tenths (small if the populations are young), so the luminosities are typical of luminous galaxies.

The brightest companion (#1) has an exponential inner luminosity profile (~ 1 kpc) but a peculiar profile at larger radii. None of the companions has any structure at near infrared wavelengths - they are smooth and symmetrical and well-resolved, even though their colors suggest young stellar populations. They have bright compact nuclei.

7. 0849+120

This QSO is faint at $v = 20.5$, and is radio-quiet. It lies in a field with several other objects with a wide range of brightness and size. Only two of these are likely to be associated with the QSO (see figure 8): the others are too bright and/or too large.

The closest companion (#1) lies about $5''$ to the N, and slightly E, and in the K band image there may be a faint connecting luminous bridge between it and the QSO. However, this bridge is not seen in H band and there are sky background fluctuations which are comparable. The QSO itself is not measurably extended.

8. 1236–003

This QSO is at $z = 2.18$, is optically moderate at $v = 19.1$, and is radio-quiet. It has three companions all at about $15''$ separation and well separated in direction. Table 8 shows their magnitudes, which are too bright to be likely associated galaxies. Their sizes and colors are also consistent with their being at considerably lower redshift - perhaps 0.2.

The closest companion to 1236–003 is seen most clearly in the H band image, at some $4''.3$ to the NW. It is also (just) detected in the K band image. There is no detectable flux connection to the QSO, or tidal elongation. In both H and K band, the QSO images appear to have extended flux to the NE, curving to the E. This is faint enough to be questionable, but suggestive as it appears identically in both H and K . While colors are subject to errors at these faint levels, the companion is blue and the extended flux is redder than the QSO, as given in Table 8.

The structure and companion seen here are fainter than in the radio-loud objects, and the field is otherwise lacking in companions as bright as seen in the radio-loud objects. However, the data are suggestive of some tidal event in 1236–003 too.

9. 0552+398

This is a bright radio-loud QSO at $z = 2.36$. It lies in a very crowded field, in which our K band image clearly shows 20 objects besides the QSO. Of these, 10 are bright or large enough that they cannot be companions. Nine of the other ten objects form a band on either side of the QSO, some $5''$ wide and at least the length of the image – $36''$. These objects are compact and in the range 4 to 6 magnitudes fainter than the QSO. The table shows their magnitudes and also H band absolute magnitudes assuming they have standard

colors ($H - K = 1$) for a young population at this redshift. We have omitted the galaxies that are too bright to be companions for brevity in the table. Figure 1 shows the positions of most of them - a few lie outside the boundary shown because of the dither. The galaxies appear more luminous than those in other fields - possibly because there are significant negative K-corrections at this higher redshift, as would be the case for young populations: as in all cases, we have adopted no k-correction. Also, $H\alpha$ is shifted into the K band at this redshift, which might increase the flux if these galaxies have Balmer emission.

The nearest companion lies about $2''.9$ to the N and there appears to be curved connecting luminosity to the QSO, suggesting tidal interaction (see figure 10). Thus, we have evidence that the QSO is a member of a dense group of compact galaxies of comparable luminosity to the host and each other. The FWHM of the companions is $0''.35$, and all are round with no structure and a bright nucleus.

The radio source is compact, luminous, variable, GHz-peaked, and has structure seen only with VLBI: a weak halo extended more E-W than N-S. This suggests the source is young.

10. 1307+297

This is a radio-quiet QSO at redshift 3.09. Our exposures are shorter than for the other objects by a factor 3 to 4 so that our detection limits are significantly different. In addition, it has a very bright guide star which gives a high level of scattered light over much of the $36''$ field. Nevertheless, we easily see one galaxy $11''$ away in H -band at $H=20.3$ and can detect it at $K=20.5$, but no other objects. The blue color and brightness indicate it is likely to be a foreground object. The QSO is measured at $H=19.1$ and $K=18.1$, so its catalogue value of $v=18.6$ seems too bright, or it is currently in a faint state.

Given the poorer detection threshold and the bright guide star, our observation of this radio-quiet object provides weak evidence that it is not in a dense galaxy environment. The QSO itself does not appear to be resolved in these exposures.

11. 0104+0215

This QSO is radio-quiet and has redshift 4.16. Because of its high redshift we exposed much longer, and mainly in K-band (rest frame U/B) on this field. The field appears empty and has one other object, of brightness similar to the QSO, that we detect with similar limits to our other fields. However, when we compare combined integrations of about 11000

secs each, from two separate nights, some very faint objects and features are seen in both. Table 10 shows the measures of these and their positions are shown in Figure 1. Figure 11 shows the environment of the QSO.

The objects are fainter than any others measured, and some are very compact, while others are larger and diffuse. The group 3, 4, 5 are all diffuse and have diameters of $2''$ to our limit of detection. The compact objects are all $0''.4$ to $0''.6$ across. At the QSO redshift they have absolute K magnitudes of ~ -25 without k -correction.

The QSO itself appears to be extended in a near E-W direction, which is almost perpendicular to the radius vector to the guide star, and so not likely to be due to PSF effects, which are radial. The extended light is stronger on the W side. In addition, there is a suggestion of a faint smooth tail to the N and curving to the E, as seen in the figure. This is so faint that it may be an artifact, and is seen more clearly on one night’s image than the other. Seeing and transparency were similar on the two nights. The shorter H -band exposure does not reveal any of these objects or features.

We will discuss the PSF removal of this (and other) objects more rigorously in a separate publication. Nevertheless, we have suggestive evidence from our K -band data that this high redshift object has an asymmetrical host galaxy, some nearby companions, and maybe a tidal tail as well. The QSO magnitude is 18.3 in H and 18.0 in K , giving it a very blue color.

12. Discussion

Our relatively small QSO sample is fairly well-matched between radio-loud and radio-quiet at redshifts near 1, 1.6, and 2.2. The small pixel size has required long integration times, so that our images are limited by sky noise. The AO camera performance has produced images with FWHM well under $0''.2$ in most bands and objects: we were careful to observe only those objects which have guide stars close and bright enough to yield optimal correction. We have reported only data obtained in good natural seeing and clear conditions. Thus, the data on all objects are very comparable.

The extended features seen in most objects are faint, so that we may be missing some fainter structure that is lost in the sky noise. This is increasingly true as we move through the J , H , K range, so that the faintest structures are seen only in J band in all 3 cases where we observed in J . It appears that there is no very small-scale structure present in any of the objects, so we binned the data 4×4 (to effective pixels $0''.14$) to decrease the sky noise and make some measurements easier.

It is notable that we find faint structures in all observed objects to redshift 2.4, and probably also in the $z=4.1$ object. These structures are smooth in overall shape and brightness, and characteristic of tidal tails of stars from a galaxy, rather than blue and knotty structure characteristic of active star formation. However, our observations are at relatively long restframe wavelengths so we are not very sensitive to star-formation activity. It will be important to obtain shorter wavelength (i.e. CCD) data of high resolution to check our conclusion that tidal activity is more important than star formation. In any event, the data suggests that tidal events are an even more marked cause (or effect) of QSO activity at higher redshifts, than the well-known observations of interactions at lower redshifts.

In our sample to date, there is also a notable difference in the galaxy environment between radio-loud and radio-quiet QSOs, with the latter generally being located in poorer environments. The exceptions are the radio-loud but apparently isolated QSO 0804+499, and possibly the radio-quiet QSO 0104+022, although the companions to the latter are detected only because of our much deeper exposure in this field. We caution against a general conclusion with our small and incomplete sample, but note that this environmental difference is an extension of what is seen at lower redshifts, although we are only able to detect the brightest of possible companion galaxies. We note also that at visible wavelengths, Hutchings (1995) finds excess galaxies around both radio-loud and radio quiet QSOs. Thus, the difference may possibly be in the population age of companions. The space density of the NIR-observed companions to the radio-loud QSOs is very high - on average the separation between galaxies in the four crowded fields observed is 35 kpc. This suggests that merging will be an important part of the activity in these groups with time. Until we can study the environments to greater depth, we cannot tell if these are simply small groups or may be the cores of what will evolve to major cluster at lower redshift, by accretion from the field. Here too, deep CCD observations with high resolution are called for, to take advantage of the lower sky brightness.

One possible related issue is the connection between objects in the field and foreground absorbers. The Hewitt Burbidge (1993) catalogue has references to absorbers in only one object: 1540+180. Thus, this is the only known case where we may have likely foreground confusion. (However, it may be that other QSOs have not been observed with sufficient resolution to detect foreground absorbers.) We have noted in the section on 1540+180 that we may be seeing absorbing objects from the $z=1.46$ system, but not the others at $z\sim 0.7$. At this point, our main inferences about the QSO environments are not affected by known foreground absorbers.

Five of our sources are radio-loud. Two have no published radio maps (but are

almost certainly compact) and two others are unresolved in the radio. The only source with comparably-sized resolved radio structure is 1540+180, which has some (but not all) corresponding visible/near infrared structure. In this connection we also note the radio-loud QSO 1345+580 (Hutchings 1998) has $L\alpha$ emission at the places of bright radio knots in a very bent radio structure.

In the only field where we have two colors, and hence can estimate redshifts, we find that there are several companions that appear to lie at the QSO redshift. In other fields, we eliminate foreground objects by their larger size and scale length, and in many cases their brightness. The remaining, possible companion galaxies are all much more compact - less than one arcsec (5 kpc) in diameter. They are mostly round and featureless except for unresolved nuclei in some. For galaxies where the signal-to-noise is sufficient to measure luminosity profiles they are neither pure disk nor power law. The small size is consistent with reports for other high redshift galaxies (e.g. Steidel et al. 1996). The smooth round morphologies suggest that the older stellar populations that we are imaging have regular shapes, in contrast with more irregular shapes seen in visible (rest UV) wavelengths, where we are seeing the clumpy nature of star-forming regions and, possibly, results of dust obscuration. The small size and relatively high luminosity of these objects suggests that they represent a stage of the early formation of galaxies. The high space density in some fields also suggests that these galaxies may merge with time, to form present day large galaxies. A similar scenario has been proposed by Rottgering(1997) for the dense environments around high redshift radio galaxies.

References

- Aretxaga I., Boyle B.J., Terlevich R.J., 1995, MNRAS, 275, 27
- Aretxaga I., Le Mignant D., Melnick J., Terlevich R.J., Boyle B.J., 1998, MNRAS, 298, L13
- Bahcall J.N., Kirhakos S., Saxe D.H., Schneider D.P., 1997, ApJ, 479, 642
- Campos A. et al 1998 1998 ApJL, in press (astro-ph/9809146)
- Casali, M. & Hawarden, T. 1992, JCMT-UKIRT Newsletter, 4, 33
- Heckman T.M., Lehnert M.D., van Breugel W., Miley G.K., 1991, ApJ, 370, 78
- Hewitt, A. & Burbidge, G. 1993, Ap. J. Suppl., 87, 451
- Hutchings J.B. & Neff S.G., 1992, AJ, 104, 1
- Hutchings J.B., Holtzman J., Sparks W.B., Morris S.C., Hanisch R.J., Mo J., 1994, ApJ, 429, L1
- Hutchings J.B. 1995, AJ, 109, 928
- Hutchings J.B. & Neff S.G. 1997a, AJ, 113, 550
- Hutchings J.B., & Neff S.G., 1997b, AJ, 113, 1514
- Hutchings J.B. 1998, AJ, 116, 20
- Hutchings J.B., Crampton D., Morris S.L., Steinbring E., 1998, PASP, 110, 374
- Lehnert M.D., Hackman T.M., Chambers K.C., Miley G.K., 1992, ApJ, 393, 68
- McLeod K. K. & Rieke G. H., 1995, ApJ, 454, L77
- Pentericci L., et al 1998, AAp, in press (astro-ph/980956)
- Rigaut, F. et al. 1998, PASP, 110, 152
- Rottgering, H. 1997, IAU Symp., 186, 184
- Steidel, C.C., Giavalisco, M., Pettini, M., Dickinson, M. & Adelberger, K.L. 1996, ApJ, 462, L17

Table 1: Table of near infrared imaging results - ordered by redshift

QSO	m_v	z	Radio	Comment
0915–213	17.5	0.85	RL	Resolved host, knot, tail, compact cluster
1055+019	20.5	1.06	RQ	Resolved, faint jet or extended companion *
0804+499	17.5	1.43	RL	~Unresolved, tail
1337–013	18.7	1.61	RL	Compact group, poss connection to nearest knot
1540+180	18	1.66	RL	Resolved, tail, compact cluster
0849+120	20.5	1.76	RQ	2 possible companions, maybe tail to one
1236–003	19.1	2.18	RQ	~Unresolved, poss tail & nearby companion
0552+398	18	2.37	RL	Interacting? dense cluster
1307+297	18.6	3.09	RQ	Short exposures, unresolved, one companion
0104+022	19.7	4.16	RQ	Resolved + tail? faint companions
* See Hutchings et al 1998				

Table 2. Journal of observations - ordered by RA

Name	Month 1998	Exposure in secs			Guide * mag R	Offset arcsec	FWHM (")		
		J	H	K			J	H	K
0104+022	Jan	–	4800	24000	13.8	14	–	0.32	0.17
0552+398	Jan	–	-	9300	11.7	26	–	–	0.24
0804+499	Mar	6000	7200	-	14.6	11	0.23	0.24	–
0849+120	Jan	–	9600	7200	13.4	16	–	0.16	0.19
0915–213	Mar	–	7200	7200	13.2	12	–	0.16	0.16
1236–003	Jan	–	9600	7200	12.8	19	–	0.25	0.15
1307+297	Jan	–	1500	1800	8.1	21	–	0.15	0.15
1337–013	Mar	7200	7800	–	13.4	20	0.19	0.15	–
1540+180	Mar	6000	6000	6000	12.6	17	0.23	0.14	0.13

Table 3. Photometry in 0915–213 field

Object	K	H	$H - K$	M_H
Q	15.9	16.7	0.8	−25.8
Q _{host}	20.3	20.6	0.3	−21.9
1	17.9	18.0	0.1	star
2	18.7	19.1	0.4	−23.4
3	21.0	21.7	0.7	−20.8
4	-	19.5	-	−23.0
5	19.1	19.8	0.7	−22.7
wisp	20.5	21.4	0.9	−21.1

Table 4. Photometry in 0804+499 field

Object	J	H	$J - H$	M_H
Q	16.9	15.9	1.0	−27.0
Q _{jet}	22.5	21.9	0.6:	−21.4
star	17.1	16.6	0.5	- - -
1	19.2	18.7	0.5	−24.7

Table 5. Photometry in 1337–013 field

Object	H	J	$J - H$	M_J
Q	17.3	18.1	0.8	−26.2
nearby knot	22.3	22.3	0.0	−21.9
1	22.5	22.4	−0.1	−22.0
2	20.4	21.2	0.8	−23.1
3		22.7		
4		22.8		

Table 6. Photometry in 1540+180 field

Object	K	H	J	$J - H$	$H - K$	M_J
Q	17.6	18.3	19.0	0.7	0.7	−25.4
Q _{host}	19.2					
1	18.7	19.6	20.3	0.7	0.9	−24.1
2	19.7	20.7	21.6	0.9	1.0	−22.8
3	19.4	20.6	21.7	1.2	1.2	−22.9
4	20.5	21.4	22.0	0.6	0.9	−22.6
5	18.5	19.0	20.0	1.0	0.5	

Table 7. Photometry in 0849+120 field

Object	H	K	$H - K$	M_H	
Q	19.5	19.1	0.4	−25.0	
1	22.2	21.3	0.9	−22.3	Compact
2	18.8	18.5	0.3		Large
3	22.2	-	-		Large faint
4	22.1	-	-	−22.4	Compact
5	19.7	19.6	0.1		Bright
6	17.5	17.3	0.3		Star

Table 8. Photometry in 1236−003 field

Object	H	K	$H - K$	M_H
Q	17.4	16.7	0.7	−27.9
1	19.4	19.2	0.2	
2	18.3	17.9	0.4	
3	20.3	19.8	0.5	
4	22.0	21.9	0.1	−23.3
QSO ext	21.0	20.0	1.0	−22.3

Table 9.
Photometry in
0552+398 field

Object	K	M_H
Q	15.2	−29.3
1	19.1	−25.4
2	20.7	−23.8
3	20.9	−23.6
4	20.5	−24.0
5	19.3	
9	19.7	−24.8
10	19.7	−24.8
13	20.4	−24.1
14	20.7	−23.8
17	20.6	−23.9
18	20.4	−24.1

Table 10.
Photometry in
0104+022 field

Object	K
Q	18.0
1	21.8
2	22.2
3 (large)	20.6
4 (large)	20.9
5 (large)	21.4
6 (star?)	18.7
7	21.7
8	21.8
9	20.9

Captions to figures

Fig. 1.— Location of detected objects in all fields. The field side is $36''$. N is up and E to the left. Some objects not shown were seen in partially exposed larger field due to dither patterns. The guide star is indicated as GS when it is in the field. The QSO is indicated by Q, assumed foreground galaxies by F, and stars by *. Photometry of most numbered galaxies is in the Tables.

Fig. 2.— K and H band images of 0915–213 (lower bright object). The images have been binned 4×4 (to $0''.14$ pixels) to enhance faint features. N is up and E to the left. Note the extension of the QSO to the NE and the nearby compact object to the NW. There is probably a faint arc of flux to the NW, seen in both bands, but more clearly in K .

Fig. 3.— J band image of 0804+499 with full resolution and with 4×4 pixel binning to enhance visibility of faint jet.

Fig. 4.— J and H band images of 1337–013, showing companion to NE and closer companion to the S, possibly with connecting flux. The greater visibility in J band may be due to redshifted [O III] emission.

Fig. 5.— 4×4 binned images in 3 bands around 1540+180. The QSO is extended to the NE, particularly in K band. In the H -band this extends to a long faint wisp terminating at a round blob, which is visible in all three bands. The radio structure to the N lies close to this near infrared wisp.

Fig. 6.— near infrared 2-color plot showing population evolution after a 1 Gyr starburst, at different redshifts. Symbols along the tracks mark the ages indicated, and the point below 1 is 0.1 Gyr. The dotted line shows the effect of extinction on the $z=1$ track. The QSO and companions 1, 2, 3, and 4 lie close to the positions expected for young populations at the QSO redshift of 1.7, possibly with some dust. Object 5 is probably a reddened foreground galaxy at $z < 0.3$.









Fig. 7.— Azimuthally averaged profiles of QSO 1540+180 and the PSF, approximately corrected for distance from the guide star.

Fig. 8.— H and K band images of the field around QSO 0849+120, binned 4×4 . In the K band there is a luminous connection to the nearest companion, to the NE.

Fig. 9.— H and K band images, binned 4×4 , of QSO 1236–003. There is a compact companion to the NW (seen more clearly in H), and connected extended flux to the NE (seen similarly in both H and K).


Fig. 10.— K band image, binned 4×4 of $4'' \times 3''$ field around QSO 0552+398. There is probably connecting flux to the compact companion to the N. Most of the other objects are thought to be compact companions to the QSO.

Fig. 11.— Deep K-band image around QSO 0104+022. The QSO is the bright object, which is extended almost E-W (the guide star lies to the south, and the lower left contains its diffraction spike). The QSO also appears to have a curved filament to the NE. Faint objects which may be companions lie at about equal distance to the W and SW.



Tickmarks are 5"


0915-213 8 X 12"



H


K

0804+499 J BAND Z=1.43





8 X 11 ARCSEC

1337-013 10 X 12"



J

H





1540+180

J H

K

11 X 12"






0849+120 12 X 8"



H

K


1236-003 11 X 11"



H


K

0552+398 K



28 X 22"

0104+022



12 X 11 ARCSEC